(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(43) International Publication Date 3 June 2004 (03.06.2004)

PCT

(10) International Publication Number WO 2004/045609 A2

- (51) International Patent Classification⁷: A61K 31/4412, A61P 17/00
- (21) International Application Number:

PCT/IB2003/005222

(22) International Filing Date:

19 November 2003 (19.11.2003)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data: MI2002A002447

19 November 2002 (19.11.2002)

(71) Applicant and

(72) Inventor: GHISALBERTI, Carlo [BR/BR]; Rua Luis Dias 85/61, ITAIM - BIBI, 04542-080 Sao Paulo (BR).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



(54) Title: HYDROXYPYRIDINONES FOR THE LOCAL TREATMENT OF SKIN MICROCIRCULATORY DISORDERS

(57) Abstract: The application of hydroxypyridonone in effective amounts as external agent for patients suffering from skin microcirculatory disorders (SMD) provides for a significant amelioration of subject conditions. Accordingly, the new use of hydroxypyridonones as external anti-inflammatory agent thereby combined with the depletion of hemosiderin residues offers a suitable treatment to SMD sufferers.